

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 064 846 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

03.01.2001 Bulletin 2001/01

(51) Int. Cl.⁷: **A01N 47/44, A01N 37/24**

(21) Application number: 99951182.7

(86) International application number:

PCT/JP99/06070

(22) Date of filing: 01.11.1999

(87) International publication number:

WO 00/25587 (11.05.2000 Gazette 2000/19)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(72) Inventors:

- URIHARA, Ichirou
Sakura-shi, Chiba 285-0817 (JP)
- TSUBOI, Hiroyuki
Yachiyo-shi, Chiba 276-0034 (JP)

(30) Priority: 02.11.1998 JP 31240698

(71) Applicants:

- DAINIPPON INK AND CHEMICALS, INC.
Itabashi-ku, Tokyo 174-8520 (JP)
- NIHON BAYER AGROCHEM K.K.
Tokyo 108-8572 (JP)
- Bayer Aktiengesellschaft
51368 Leverkusen (DE)


(74) Representative:

Albrecht, Thomas, Dr. et al
Kraus, Weisert & Partner
Patent- und Rechtsanwälte
Thomas-Wimmer-Ring 15
80539 München (DE)

(54) **BACTERIOCIDIC COMPOSITIONS FOR AGRICULTURAL AND HORTICULTURAL USE**

(57) The novel agricultural and horticultural fungicidal compositions containing as active ingredients a combination of

A) 1, 1'-iminiodi(octamethylene)diguanidine of the formula




(iminooctadine)

or an acid addition salt thereof and

B) 1,1'-iminiodi(octamethylene)diguanidine of the formula

EP 1 064 846 A1



(II)

(fenhexamid)

are outstandingly active as fungicides.

Description

TECHNICAL FIELD

[0001] The present invention relates to novel agricultural and horticultural compositions having fungicidal properties. The compositions contain the known iminoctadine and acid addition salts thereof on the one hand and the known N-(2,3-dichloro-4-hydroxy-phenyl)-1-methyl-cyclohexane carboxamide on the other hand.

BACKGROUND ART

[0002] It has already been known that iminoctadine and acid addition salts thereof are effective against a broad range of diseases that occur in important crop plants (see JP-A Sho 62-209003). Further, it has been described already that N-(2,3-dichloro-4-hydroxyphenyl)-1-methyl-cyclohexane carboxamide has a good fungicidal activity and is effective against gray mold in grapes and citrus fruits as well as against brown rot in peaches (see JP-A Hei 2-11551). However, the activity of these compounds is not always fully satisfactory for the control of severe infections with gray mold. Moreover, the control of gray mold fungus has become an even more serious problem due to the emergence of strains which are resistant to benzimidazoles, dicarboximides and diethofencarb


DISCLOSURE OF INVENTION

[0003] The present invention provides new compositions, which are excellently suitable for the control of phytopathogenic fungi, such as not only gray mold, but also fungal strains which are resistant to the above-mentioned agents. Furthermore, the compositions according to the invention are effective against a broad range of plant diseases even if the infection is severe.

[0004] The present invention has been completed after intensive research to resolve the above-mentioned problems. It has been found that compositions containing iminoctadine or acid addition salts thereof and N-(2,3-dichloro-4-hydroxy-phenyl)-1-methyl-cyclohexane carboxamide are very effective against fungi causing disease in plants, including not only gray mold, but also strains, which are resistant to the above-mentioned agents. It has also been found that the compositions according to the invention exert an excellent control effect even if the infection is severe.

[0005] It has been found now that the novel agricultural and horticultural fungicidal compositions containing as active ingredients a combination of


A) 1,1'-iminiodi(octamethylene)diguanidine of the formula



(iminoctadine)

or an acid addition salt thereof and

B) N-(2,3-dichloro-4-hydroxy-phenyl)-1-methyl-cyclohexanecarboxamide of the formula



(fenhexamid)

exhibit a very good fungicidal activity.

[0006] Surprisingly, the fungicidal effect of the active compound combinations according to the invention is substantially greater than the sum of the effects of the individual active compounds. It is, therefore, a matter of an unforeseeable, genuine synergistic effect and not simply of an addition to the effect.



BEST MODE FOR CARRYING OUT THE INVENTION

[0007] The 1,1'-iminiodi(octamethylene)diguanidine is characterized by the formula (I) and is known under the common name iminotadine.

[0008] Acid addition salts of iminotadine include carboxylates, sulfonates, hydrochloride, carbonate, lauryl sulfate, sulfate, phosphate and the like. Examples of carboxylates include acetate, oxalate, laurylate, oleate, citrate and the like. Examples of sulfonates include alkyl sulfonate, alkyl benzene sulfonate, and the like. Examples of alkyl sulfonates include lauryl sulfonate, dodecyl sulfonate and the like.

[0009] Examples of alkyl benzene sulfonates include dodecyl benzene sulfonate, p-toluene sulfonate and the like. Preferred acid addition salts are carboxylates and sulfonates, such as alkyl benzene sulfonates. Particularly preferred are alkyl benzene sulfonates having 1 to 18 carbon atoms, specifically 10 to 13 carbon atoms, in the alkyl group.

[0010] The 1,1'-iminiodi(octamethylene)diguanidinium tris(alkyl-benzenesulfonate) [iminotadine-tris(albesilate)] may be mentioned as an example of an acid addition salt of iminotadine. This compound can be represented by the following formula



$n = 10 \text{ to } 13.$

[iminotadine tris(albesilate)].

[0011] The iminotadine of the formula (I) and acid addition salts thereof are known (see JP-A Sho 62-209003 and "The Pesticide Manual", 11th Edition 1997, pages 709-712).

[0012] The N-(2,3-dichloro-4-hydroxy-phenyl)-1-methyl-cyclohexane carboxamide of the formula (II) is also a known compound (see JP-A Hei 2-11551 and EP-A 0339418).

[0013] The synergistic effect is particularly clear when the active compounds are present in the compositions according to the invention in particular weight ratios. However, the weight ratios of the active compounds can be varied

within a relatively wide range. In general, from 0.5 to 10 parts by weight, preferably from 1 to 5 parts by weight, of the compound of the formula (II) are allotted to 1 part by weight of a compound of group (A).

[0014] The agricultural and horticultural fungicidal compositions according to the invention possess very good fungicidal properties and can be used for the control of phytopathogenic fungi in agriculture and horticulture. They are suitable for the control of fungi, which are resistant to some other chemical compounds, and against fungi causing plant diseases, such as late blight, gray mold, leaf mold, leaf spot, anthracnose, and early blight in tomatoes; downy mildew, powdery mildew, gray mold anthracnose, gummy stem blight, phytophthora rot, scab, and *Corynespora* leaf spot in cucumbers; stem blight, leaf spot and gray mold in asparagus; anthracnose and gray mold in strawberries; powdery mildew, anthracnose, and gummy stem blight in pumpkins; *Sclerotinia* rot in cabbages; anthracnose in garland chrysanthemums; powdery mildew, *Sclerotinia* rot, anthracnose and gummy stem blight in watermelons; *alternaria* leaf spot, gray mold, and gray mold neck rot in onions; powdery mildew, gray mold and black rot in eggplants; *alternaria* leaf spot in welsh onions; leaf spot and *alternaria* leaf spot in Chinese cabbages; *Sclerotinia* rot and gray mold in peppers; powdery mildew, anthracnose, and gummy stem blight in melons; gray mold in lettuce; scab and gray mold in Japanese apricots; powdery mildew, anthracnose, gray mold and leaf spot in Japanese persimmons; gray mold and scab in citrus fruits; scab, black spot and *physalospora* canker in Japanese pears; gray mold, ripe rot, leaf spot, anthracnose, gummy stem blight, black rot, and swelling arm in grapes; scab, brown rot, *Phomopsis* rot, and anthracnose in peaches; scab, *alternaria* leaf spot/ring rot, blotch, sooty blotch, fly speck, fruit spot, and blossom blight in apples; stalk blight anthracnose, gray mold and *Sclerotinia* rot in Azuki beans; anthracnose, gray mold, and *Sclerotinia* rot in kidney beans; ring spot in broad beans; purple speck in soybeans; powdery mildew in tobacco; *Romalaria* leaf spot and *cercospora* leaf spot in sugar beets; anthracnose, gray blight, shoot blight and gray mold in tea; and black spot, anthracnose, gray mold, and powdery mildew in flowers. In particular, the agricultural and horticultural fungicidal compositions according to the invention are suitable for the control of fungi, which are resistant to some other chemical compounds, and against fungi causing plant diseases, such as gray mold in tomatoes, cucumbers, eggplants, and strawberries, or gray mold in fruits such as grapes, peaches, and citrus fruits.

[0015] The agricultural and horticultural fungicidal compositions can be used during the growing period of crops such as asparagus, strawberries, pumpkins, cabbages, cucumbers, garland chrysanthemums, watermelons, onions, tomatoes, eggplants, green onion, Chinese cabbages, peppers, spinach, Oriental melons, melons, lettuces, Japanese apricots, Japanese persimmons, citrus fruits, Japanese pears, grapes peaches, apples, Azuki beans, kidney beans, broad beans, soybeans, tobacco, sugar beets, tea, and flowers (i.e., roses, chrysanthemums, carnations, cyclamen, gentian, etc.). For grapes in particular, the agricultural and horticultural fungicidal composition can be preferably used between the stage prior to flowering and the young fruit stage in their growing period.

[0016] The agricultural and horticultural fungicidal compositions according to the invention can be used as such or in the form of customary formulations, such as wettable powders, suspension concentrates, dustable powders, emulsifiable concentrates, oil miscible liquids, tablets, granules, aerosols and the like. The formulations can be prepared by customary methods, for instance by mixing the active compounds with other components, such as extenders, surfactant agents, other additives, or suitable carriers, including water. The agricultural and horticultural fungicidal compositions according to the invention are preferably used in the form of wettable powders or suspension concentrates.

[0017] Examples of extenders include kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite, diatomaceous earth, calcium carbonate, hydrated lime, quartz sand, ammonium sulfate, carbamide, vermiculite, wood chips, starch, clay, talc, bentonite, white carbon, etc., in the case of solid carriers, arid alcohols, kerosene, naphtha, aromatic compounds (xylene, toluene, phenol, etc.), alkyl naphthalene, cyclohexane, benzene, acetone, N-methyl pyrrolidone and the like in the case of liquid carriers.

[0018] Examples of surfactant agents include polyoxysorbitan monooleate, ethylene oxide/propylene oxide copolymers, lignin sulfonate, sorbitan ester, soaps, alkyl benzene sulfonate, fatty acid amine salts, quaternary ammonium salts, alkyl pyridinium salts, alkyl aminoethyl glucine, alkyl dimethyl betaine, polyglycol sulfate ester, alkyl amino sulfonate, isopropyl phosphate, carboxymethyl cellulose, hydroxypropyl cellulose, xanthane gum, diaryl alkyl disulfonate, alcohol sulfate, alkyl aryl sulfonate, polyoxyethylene alkyl aryl ether, polyoxyethylene glycol ether, polyoxyethylene sorbitan monoalkylate, and the like.

[0019] Other additives include adjuvants like carboxymethyl cellulose, ethylene glycol and propylene glycol, as well as additives typically used during formulation.

[0020] The formulations generally contain between 30 and 90 per cent by weight of the active compounds, preferably between 50 and 80 per cent by weight. The active compounds can be present in the formulations in admixture with other known active compounds, such as fungicides, insecticides, acaricides, nematocides, fertilizers and plant-growth regulators.

[0021] The agricultural and horticultural fungicidal composition according to the invention are preferably used in the form of dispersions, which can be obtained by diluting the formulations with water. It is also possible to add other fungicides or pesticides to the formulations and then to disperse them. The concentration of the active compounds in the dispersions can be varied according to the period of use, method, site, form of the agent, targeted disease and targeted crop. However, the range for the acid addition salt of the iminoctadine is between 25 and 24,000 ppm, preferably

between 60 and 200 ppm (when using iminoctadine, between 7.8 and 7,500 ppm, preferably between 19 and 63 ppm), while the range for fenhexamid is between 100 and 40,000 ppm, preferably between 150 and 300 ppm. In particular, in the case where the composition is used for fruits such as grapes and peaches, the concentration of the active compounds in the dispersions is such that the range for the acetic acid salt of the iminoctadine is between 25 ppm and 24,000 ppm, preferably between 200 ppm and 1,000 ppm (when using iminoctadine, between 7.8 ppm and 7,500 ppm, preferably between 62 ppm and 313 ppm), while the range for the fenhexamid is between 100 ppm and 40,000 ppm, preferably between 250 ppm and 1,500 ppm.

[0022] The dispersing amount will vary depending on the period of use, method, site, form of the agent, target disease, target crop, etc. For example, when spraying over vegetables such as tomatoes and cucumbers, the amount of the agent in the concentration of the effective component that is dispersed is preferably between 150 and 300 L per 10 are, while when spraying over fruits such as grapes and peaches, the amount of agent in the concentration of the effective component that is dispersed is preferably between 200 and 700 L per 10 are.

Examples

[0023] The present invention is illustrated by the following examples and use examples without being limited thereto. The term "part", as used below, means "parts by weight".

Example 1 (Wettable powder 1)

[0024]

iminoctadine tris(albesilate)	200 parts
fenhexamid	300 parts
clay	210 parts
lignin sodium sulfonate	20 parts
polyoxyethylene alkyl phenyl ether	50 parts
polyvinyl alcohol	20 parts
white carbon	200 parts
	1000 parts

[0025] The above-mentioned components are placed in a ribbon mixer, mixed well, and were pulverized sufficiently in an atomizer to obtain a wettable powder.

Example 2 (Wettable powder 2)

[0026]

iminoctadine tris(albesilate)	120 parts
fenhexamid	200 parts
clay	470 parts
lignin sodium sulfonate	20 parts
polyoxyethylene alkyl phenyl ether	50 parts
polyvinyl alcohol	20 parts
white carbon	120 parts
	1000 parts

EP 1 064 846 A1

[0027] The above-mentioned components are placed in a ribbon mixer, mixed well, and were pulverized sufficiently in an atomizer to obtain a wettable powder.

Example 3 (Wettable powder 3)

[0028]

iminotadine tris (albesilate)	40 parts
fenhexamid	400 parts
clay	430 parts
lignin sodium sulfonate	20 parts
polyoxyethylene alkyl phenyl ether	50 parts
polyvinyl alcohol	20 parts
white carbon	40 parts
	1000 parts

[0029] The above-mentioned components are placed in a ribbon mixer, mixed well, and were pulverized sufficiently in an atomizer to obtain a wettable powder.

Example 4 (Wettable powder 4)

[0030]

iminotadine tris(albesilate)	200 parts
fenhexamid	100 parts
clay	310 parts
lignin sodium sulfonate	20 parts
polyoxyethylene alkyl phenyl ether	50 parts
polyvinyl alcohol	20 parts
white carbon	300 parts
	1000 parts

[0031] The above-mentioned components are placed in a ribbon mixer, mixed well and were pulverized sufficiently in an atomizer to obtain a wettable powder.

Example 5 (Suspension concentrate 1)

[0032]

A)	iminotadine tris(albesilate)	120 parts
	polyoxyethylene lauryl ether	30 parts
	sorbitan oleic acid ester	30 parts
	propylene glycol	50 parts
	water	290 parts
B)		520 parts
	fenhexamid	200 parts
	polyoxyethylene styryl phenyl ether	30 parts
	water	250 parts
		480 parts

[0033] A) was mixed well in a homogenizer and dispersed, while B) was wet-ground in a bead mill. A) and B) were then thoroughly mixed together to form a suspension concentrate.

Comparative agent 1 (iminotadine tris(albesilate) wettable powder)

[0034]

iminotadine tris(albesilate)	400 parts
polyoxyethylene lauryl ether	50 parts
clay	150 parts
white carbon	400 parts
	1000 parts

[0035] The above-mentioned components were placed in a ribbon mixer, mixed well, and were pulverized thoroughly with an atomizer to obtain a wettable powder.

Comparative agent 2 (fenhexamid wettable powder)

[0036] Fenhexamid wettable powder (effective component 50% by weight) manufactured by Japan Bayer Agrochem.

Comparative agent 3

[0037] 3-(3,5-dichlorophenyl)-N-isopropyl 2,4-dioximidazolidine-1-carboxamide (hereinafter referred to as IPRIDI-ONE) wettable powder (effective component 50% by weight) manufactured by Nissan Chemical Industry Co.

Comparative agent 4

[0038] N-(4-methyl-6-prop-1-ynylpyrimidin-2-yl)aniline (hereinafter referred to as MEPANIPYRIM) wettable powder (effective component 40% by weight) manufactured by Kumiai Chemical Industry Co.

Use Examples

Example A

5 Test of control effect on gray mold in cucumbers

[0039] 18 cucumber plants (commercial name: ANKORU 8) cultivated under ordinary conditions at room temperature were assigned for each agent. Tangerines inoculated with gray mold fungi (resistant/sensitive strain: benzimidazole resistant, dicarboximide sensitive) provided by Japan Plant Protection Association were suspended to serve as inoculation sources. Each of the agents was diluted with water so that the active compounds reached the values shown in the table. Cucumbers at the 10-leaf stage at the start of the experiment were sprayed a total of 4 times at one week intervals using a compressed air sprayer to disperse a sufficient amount (300L/10a) of the diluted solutions of each of the agents. Prior to each spraying, the number of harvested fruit and the number of infected fruit were noted. One week after the last spraying, all fruit was assessed for the infected fruit, and a value was obtained for the occurrence rate of diseased fruit. Control values (%) were calculated using Equation 1 below. These results are shown in Table 1.

$$\text{control value} = \frac{\left(\frac{\text{disease occurrence rate when untreated} - \text{disease occurrence rate with single or combined agent}}{\text{disease occurrence rate when untreated}} \right) \times 100}{\text{disease occurrence rate when untreated}} \quad \text{Equation 1}$$

20

[0040] The additive effect obtained when two types of agents are mixed together can be expressed as the control value's expected value (%) (hereinafter, referred to as PE) indicated in Equation 2. Therefore, it is possible to determine the presence or absence of a "synergistic effect" from the relationship with the actual control value for the mixed agent.

25

$$\text{Control value's expected value} = (m + n) - \frac{mn}{100} \quad \text{Equation 2}$$

[0041] m and n in the above equation indicate the control values for iminoctadine tris(albesilate) and fenhexamid respectively.

[0042] The control value's expected values are shown inside parenthesis under the control value column in Table 1. Determination of the synergistic effect was made based on the following standards.

PC > PE : synergistic effect

PC = PE : additive effect

35 PC < PE : antagonistic effect.

Table 1

Sample Agent	Effective component and its concentration (ppm)		Disease occurrence rate (%)	Control value
wetable powder 1	iminocadine tris(albesilate)	200 ppm	1.3	94.6
	fenhexamid	300 ppm		(84.5)
comparative agent 1	iminocadine tris(albesilate)	200 ppm	7.9	67.1
comparative agent 2	fenhexamid	300 ppm	11.3	52.9
comparative agent 4	MEPANIPYRIM	200 ppm	10.7	55.4
untreated			24.0	

50

[0043] The control value for gray mold in cucumbers for wettable powder 1 was superior to comparative agent 1 and 2 which were used separately, and was also better than conventional comparative agent 4. The control value for gray mold in cucumbers for wettable powder 1 was higher than the control value's expected value obtained from the control values of comparative agents 1 and 2, thus confirming the synergistic effect of the agricultural and horticultural fungicidal composition according to the invention.

Example B

Test of control effect on gray mold in tomatoes

5 [0044] 18 tomato plants (commercial name: MOMOTARO) cultivated under ordinary conditions at room temperature were assigned to each agent. Tangerines inoculated with the same gray mold fungi as in Example A (R/S Strain) were suspended to serve as inoculation sources. Each of the agents was diluted with water so that the active compounds achieved the values shown in the table. Tomatoes at the 10 leaf stage at the start of the experiment were sprayed a total of 5 times at one week intervals using a shoulder-mounted compressed air sprayer to disperse a sufficient amount (476L/10a) of the diluted solutions of each of the agent. Prior to each spraying, the number of harvested fruit and the number of infected fruit was noted. One week after the last spraying, the total number of fruit was assessed for the infected fruit, and a value was obtained for disease occurrence rate. Control values were calculated using Equation 1 above. The results are shown in Table 2. The control value's expected values were determined using Equation 2, and are shown inside parenthesis under the control value column in Table 2.

Table 2

Sample Agent	Effective component and its concentration (ppm)		Disease occurrence rate(%)	Control value
wetable powder 2	iminoctadine tris(albesilate)	120 ppm	1.5	93.7 (84.5)
	fenhexamid	200 ppm		
Suspension concentrate 1	iminoctadine tris(albesilate)	120 ppm	1.1	95.5 (84.5)
	fenhexamid	200 ppm		
comparative agent 1	iminoctadine tris(albesilate)	120 ppm	8.5	64.4
comparative agent 2	fenhexamid	200 ppm	10.4	56.5
comparative agent 3	IPRODIONE	333 ppm	14.4	39.7
untreated			23.9	

35 [0045] The control values for gray mold in tomatoes for wettable powder 2 and suspension concentrate 1 were superior to that of comparative agents 1 and 2 which were used separately, and was also better than conventional comparative agent. The control value for gray mold in tomatoes for wettable powder 2 and suspension concentrate 1 was higher than the control value's expected value obtained from the control values of comparative agents 1 and 2, thus confirming the synergistic effect of the agricultural and horticultural fungicidal composition according to the invention.

40 Example C

Test of control effect on gray mold in eggplants

45 [0046] 9 eggplants (commercial name: SENRYO No. 2) cultivated under ordinary conditions at room temperature were assigned to each agent. Tangerines inoculated with the same gray mold fungi as in Example A (R/S stain) were suspended to serve as inoculation sources. Each of the agents was diluted with water so that the active compounds achieved the values shown in the table. Eggplants at the 8-leaf stage at the start of the experiment were sprayed of a total of 5 times at one week intervals using a compressed air sprayer to disperse a sufficient amount (300L/10a) of the diluted solutions of each of the agents. Prior to each spraying, the number of harvested fruit and the number of infected fruit was noted. One week after the last spraying, the total number of fruit was assessed for the infected fruit, and a value was obtained for disease occurrence rate. Control values were calculated using Equation 1 below. The results are shown in Table 3. The control value's expected values were determined using Equation 2, and are shown inside parenthesis under the control value column in Table 3.

Table 3

Sample Agent	Effective component and its concentration (ppm)		Disease occurrence rate(%)	Control value
wettable powder 1	iminotadine tris(albesilate)	133 ppm	12.5	80.5
	fenhexamid	200 ppm		(67.3)
comparative agent 1	iminotadine tris(albesilate)	133 ppm	33.6	47.6
comparative agent 2	fenhexamid	200 ppm	40.0	37.6
wettable powder 3	iminotadine tris(albesilate)	27 ppm	18.5	71.1
	fenhexamid	267 ppm		(64.8)
comparative agent 1	iminotadine tris(albesilate)	27 ppm	44.8	30.1
comparative agent 2	fenhexamid	267 ppm	32.3	49.6
wettable powder 4	iminotadine tris(albesilate)	200 ppm	16.8	73.8
	fenhexamid	100 ppm		(68.1)
comparative agent 1	iminotadine tris(albesilate)	200 ppm	25.6	60.1
comparative agent 2	fenhexamid	100 ppm	51.2	20.1
comparative agent 3	IPRODIONE	333 ppm	35.0	45.4
untreated			64.1	

[0047] The effect of the comparative agent was not very high when the infection was severe. In contrast, the control value of wettable powders 1, 3, and 4 on gray mold in eggplants was superior to that of comparative agents 1-3. The control value of wettable powders 1, 3 and 4 on gray mold in eggplants was higher than the control value's expected value obtained from the control values of comparative agents 1 and 2, thus confirming the synergistic effect of the agricultural and horticultural fungicidal compositions according to the invention.


INDUSTRIAL APPLICABILITY

[0048] As explained above, agricultural and horticultural fungicidal compositions containing iminotadine or acid addition salts thereof and fenhexamid are very effective in controlling phytopathogenic fungi, such as gray mold and fungal strains which are resistant to chemical compounds.

Claims


1. Agricultural and horticultural fungicidal compositions, characterized in that they contain as active ingredients a combination of

A) 1,1'-iminiodi(octamethylene)diguandine of the formula





or an acid addition salt thereof and

B) N-(2,3-dichloro-4-hydroxy-phenyl)-1-methyl-cyclohexane carboxamide of the formula



- 10
2. Compositions according to claim 1, wherein the compound of group (A) is 1,1'-iminiodi(octamethylene)diguanidinium ris(alkyl-benzenesulfonate) of the formula
- 15



30 n = 10 to 13

35 [iminooctadine tris (albesilate)]

- 40 3. Compositions according to claim 1 characterized in that the weight ratio of active compounds of group (A) to the active compound of the formula (II) is between 1:0.5 and 1:10.
4. Method for the control of fungi, characterized in that active compound compositions according to claim 1 are applied to the fungi and/or to their habitat.
- 45 5. Use of compositions according to claim 1 for the control of fungi.
6. Process for the preparation of fungicidal compositions, characterized in that active compound combinations according to claim 1 are mixed with extenders and or surface-active agents.
- 50

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/06070

A. CLASSIFICATION OF SUBJECT MATTER

IPC.C1⁷ A01N47/44, A01N37/24

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC.C1⁷ A01N47/44, A01N37/24

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP, 70792, A1 (BAYER AG), 24 April, 1996 (24.04.96) & JP, 8-208406, A & CN, 1132591, A & AU, 699963, B & US, 5962518, A	1, 3-6
Y	US, 5728734, A (Dainippon Ink and Chemicals, Inc.), 17 March, 1998 (17.03.98) & JP, 8-225402, A	2
Y	JP, 8-113505, A (DAINIPPON INK AND CHEMICALS, INC.), 07 May, 1996 (07.05.96) (Family: none)	2
A	EP, 626135, A2 (BAYER AG), 30 November, 1994 (30.11.94) & JP, 6-329505, A & CN, 1094221, A & US, 5532262, A & US, 5672619, A & US, 5922762, A	1-6

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not

considered to be of particular relevance

"E" earlier document but published on or after the international filing

date

"L" document which may throw doubts on priority claim(s) or which is

cited to establish the publication date of another citation or other

special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other

means

"P" document published prior to the international filing date but later

than the priority date claimed

"T" later document published after the international filing date or

priority date and not in conflict with the application but cited to

understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be

considered novel or cannot be considered to involve an inventive

step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be

considered to involve an inventive step when the document is

combined with one or more other such documents, such

combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search
25 January, 2000 (25.01.00)Date of mailing of the international search report
15 February, 2000 (15.02.00)Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)